


## CLAIMS

1. Medium for detecting and/or identifying microorganisms present in a sample, comprising a culture medium and at least one substrate that can be hydrolysed to a labelled product by at least a first enzyme not free in the sample, and specific for said microorganisms, characterized in that it also comprises at least one inhibitor of at least a second enzyme, different from the first enzyme or identical to it, but free in said sample and not originating from a microorganism.
2. Detection and/or identification medium according to Claim 1, characterized in that the microorganism is a bacterium.
3. Detection and/or identification medium according to Claim 2, characterized in that said bacterium belongs to the *Salmonella* genus.
4. Detection and/or identification medium according to Claim 1, characterized in that the microorganism is a yeast.
5. Detection and/or identification medium according to Claim 4, characterized in that said yeast belongs to the *Candida* genus.
6. Detection and/or identification medium according to any one of Claims 1 to 5, characterized in that said first enzyme is an esterase.
7. Detection and/or identification medium according to Claim 6, characterized in that the inhibitor is a compound of formula (I)



in which R<sub>1</sub> is a hydrogen atom, or an alkyl, aryl or halogen group,  
R<sub>2</sub> is a hydrogen atom, or an alkyl, aryl or halogen group,  
R<sub>3</sub> is nothing, or an alkyl, aryl or NO<sub>2</sub> group.

5

8. Detection and/or identification medium according to Claim 7, characterized in that the inhibitor is O,O-diethyl p-nitrophenyl phosphate and/or O,O-dimethyl p-nitrophenyl phosphate and/or O,O-di-(2-chloroethyl)-O-(3-chloro-4-methylcoumarin-7-yl) phosphate and/or at least one derivative of these molecules.

10

9. Detection and/or identification medium according to Claim 8, characterized in that the concentration of O,O-diethyl p-nitrophenyl phosphate or its derivative in the detection medium is between 0.1 and 15 mg/l, preferably between 1 and 10 mg/l.

15

10. Detection and/or identification medium according to Claim 8, characterized in that the concentration of O,O-dimethyl p-nitrophenyl phosphate or its derivative in the detection medium is between 0.1 and 100 mg/l, preferably between 10 and 50 mg/l.

20


11. Detection and/or identification medium according to Claim 8, characterized in that the concentration of O,O-di-(2-chloroethyl)-O-(3-chloro-4-methylcoumarin-7-yl) phosphate or its derivative in the detection medium is between 1 and 1000 mg/l, preferably between 30 and 100 mg/l.

25

12. Detection and/or identification medium according to any one of Claims 1 to 5, characterized in that said first enzyme is an osidase, preferably a glucosidase.

30

13. Detection and/or identification medium according to any one of Claims 12, characterized in that the inhibitor is a compound of formula (II):




(II)

or a derivative of this compound.

14. Detection and/or identification medium according to Claim 13, characterized in that the concentration of compound of formula (II) or its derivative in the detection medium is preferably between 1 and 10 g/l, and even more preferably between 2 and 8 g/l.
15. Detection and/or identification medium according to any one of Claims 1 to 14, characterized in that said substrate is a chromogenic substrate, preferably an ester of indoxyl or of its derivatives.
16. Method for detecting and/or identifying microorganisms, comprising:

  - seeding the microorganisms to be identified onto a detection medium, according to any one of Claims 1 to 15,
  - incubating the detection medium seeded with the microorganisms to be identified, and
  - determining the presence of microorganisms by detecting the substrate(s) hydrolysed to a labelled product.
17. Use of the detection and/or identification medium according to any one of Claims 1 to 15, for identifying microorganisms.
18. Use of a compound of formula (I)




in which  $R_1$  is a hydrogen atom, or an alkyl, aryl or halogen group,

$R_2$  is a hydrogen atom, or an alkyl, aryl or halogen group,

5  $R_3$  is nothing, or an alkyl, aryl or  $\text{NO}_2$  group,  
for inhibiting a free enzyme in a sample.

19. Use of a compound of formula (II)



10 for inhibiting a free enzyme in a sample.